

RENOVO Fungicide

GROUP

1

FUNGICIDE

**COMMERCIAL
Suspension Concentrate**

ACTIVE INGREDIENT:

thiophanate-methyl

500 g/L

Contains 5-chloro-2-methyl-4-isothiazolin-3-one at 0.00113% and 2-methyl-4-isothiazolin-3-one at 0.00037 % as preservatives.

OR

Contains 5-chloro-2-methyl-4-isothiazolin-3-one at 0.001695% and 2-methyl-4-isothiazolin-3-one at 0.000555%, as preservatives.

READ THE LABEL AND ATTACHED BOOKLET BEFORE USING

REGISTRATION NO.:32097 *PEST CONTROL PRODUCTS ACT*

WARNING – POISON



Warning, contains the allergen soy.

Registrant:
Nippon Soda Co., Ltd
Shin-Ohtemachi Bldg., 2-1, 2-Chrome Ohtemachi, Chiyoda-Ku
Tokyo, Japan, 100-8165
USA Contact Number: (212) 490-0350

Canadian Representative:
Nailor Regulatory Consulting Inc.
935 Rosinke Drive, Kincardine, Ontario
N2Z 2X3
(519) 829-5422

Net Contents: 1 L to 1060 L

NOTICE TO USER:

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the **PEST CONTROL PRODUCTS ACT** to use this product in a way that is inconsistent with the directions on the label.

FIRST AID:

If swallowed: Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

TOXICOLOGICAL INFORMATION:

No specific antidote is available. Treat symptomatically.

PRECAUTIONS:

KEEP OUT OF REACH OF CHILDREN. May be harmful if swallowed or inhaled. Avoid contact with eyes, skin or clothing. Avoid breathing of dust or spray mist. Wash thoroughly with soap and water after handling. Wash contaminated clothes before re-use. Do not contaminate food or feed.

Apply only when the potential for drift beyond the area to be treated is minimal. Take into consideration wind speed, wind direction, temperature inversions, application equipment, and sprayer settings.

STORAGE:

Store this product away from food or feed.

DISPOSAL:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Follow provincial instructions for any required additional cleaning of the container prior to its disposal.
3. Make the empty container unsuitable for further use.
4. Dispose of the container in accordance with provincial requirements.
5. For information on the disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

IN CASE OF EMERGENCY INVOLVING A POISONING OR MAJOR SPILL CALL 1-866-336-2983

GROUP

1

FUNGICIDE

RENOVO Fungicide**COMMERCIAL
Suspension Concentrate****ACTIVE INGREDIENT:**

thiophanate-methyl

500 g/L

Contains 5-chloro-2-methyl-4-isothiazolin-3-one at 0.00113% and 2-methyl-4-isothiazolin-3-one at 0.00037 % as preservatives.

OR

Contains 5-chloro-2-methyl-4-isothiazolin-3-one at 0.001695% and 2-methyl-4-isothiazolin-3-one at 0.000555%, as preservatives.

READ THE BOOKLET BEFORE USINGREGISTRATION NO.: 32097 *PEST CONTROL PRODUCTS ACT*

WARNING – POISON



Warning, contains the allergen soy.

Registrant:

Nippon Soda Co., Ltd

Shin-Ohtemachi Bldg., 2-1, 2-Chrome Ohtemachi, Chiyoda-Ku

Tokyo, Japan, 100-8165

USA Contact Number: (212) 490-0350

Canadian Representative:

Nailor Regulatory Consulting Inc.

935 Rosinke Drive, Kincardine, Ontario

N2Z 2X3

(519) 829-5422

GENERAL INFORMATION:

RENOVO Fungicide is a systemic fungicide for control of certain fungus diseases of turf.

NOTICE TO USER:

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the **PEST CONTROL PRODUCTS ACT** to use this product in a way that is inconsistent with the directions on the label.

FIRST AID:

If swallowed: Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

TOXICOLOGICAL INFORMATION:

No specific antidote is available. Treat symptomatically.

PRECAUTIONS:

KEEP OUT OF REACH OF CHILDREN. May be harmful if swallowed or inhaled. Avoid contact with eyes, skin or clothing. Avoid breathing of dust or spray mist. Wash thoroughly with soap and water after handling. Wash contaminated clothes before re-use. Do not contaminate food or feed.

Apply only when the potential for drift beyond the area to be treated is minimal. Take into consideration wind speed, wind direction, temperature inversions, application equipment, and sprayer settings.

DO NOT enter or allow worker entry into treated areas to perform post-application activities during the intervals specified in the following table.

Crop	Post-Application Activity	REI and/or PHI
Sod farms	All activities	12 hours
Golf courses	All activities	Until sprays have dried

After handling RENOVO Fungicide, wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. No food, drink or tobacco should be allowed in areas of chemical storage or use. Do not apply this product in a way that will contact workers or other persons. Only protected handlers may be in the area during application.

Personal Protective Equipment (PPE):

Wear coveralls over a long-sleeved shirt, long pants, chemical-resistant gloves, shoes and socks during mixing, loading, application, clean-up and repair. Gloves are not required during application within a closed cab.

If mixing and loading more than 520 L in a day, a closed mixing/loading system is required.

For groundboom application, if applying more than 520 L in a day, a closed cab tractor is required.

Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. Remove Personal Protective Equipment (PPE) immediately after handling this product. Wash the outside of the gloves before removing. As soon as possible, wash thoroughly and change into clean clothing. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

ENVIRONMENTAL PRECAUTIONS

Toxic to aquatic organisms. Observe buffer zones specified under DIRECTIONS FOR USE.

Toxic to birds and small wild mammals.

Toxic to earthworms.

Toxic to bees. Minimize spray drift to reduce harmful effects on bees in habitats close to the application site. Avoid applications when bees are foraging in the treatment area in ground cover containing blooming weeds. To further minimize exposure to pollinators, refer to the complete guidance "Protecting Pollinators during Pesticide Spraying – Best Management Practices" on the Health Canada website (www.canada.ca/pollinators). Follow crop specific directions for application timing.

To reduce runoff from treated areas into aquatic habitats avoid application to areas with a moderate to steep slope, compacted soil, or clay.

Avoid application when heavy rain is forecast.

Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative filter strip between the treated area and the edge of the water body.

STORAGE:

Store this product away from food or feed.

DISPOSAL:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Follow provincial instructions for any required additional cleaning of the container prior to its disposal.
3. Make the empty container unsuitable for further use.
4. Dispose of the container in accordance with provincial requirements.
5. For information on the disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

IN CASE OF EMERGENCY INVOLVING A POISONING OR MAJOR SPILL CALL 1-866-336-2983

RESISTANCE MANAGEMENT RECOMMENDATIONS:

For resistance management, please note that RENOVO Fungicide contains a Group 1 fungicide. Any fungal population may contain individuals naturally resistant to RENOVO Fungicide and other Group 1 fungicides. A gradual or total loss of pest control may occur over time if these fungicides are used repeatedly in the same fields. Other resistance mechanisms that are not linked to site of action but specific for individual chemicals, such as enhanced metabolism, may also exist. Appropriate resistance-management strategies should be followed.

To delay fungicide resistance:

- Where possible, rotate the use of RENOVO Fungicide or other Group 1 fungicides with different groups that control the same pathogens.
- Use tank mixtures with fungicides from a different group that is effective on the target pathogen when such use is permitted.

- Fungicide use should be based on an Integrated Pest Management program that includes scouting, historical information related to pesticide use and crop rotation and considers host plant resistance, impact of environmental conditions on disease development, disease thresholds, as well as cultural, biological and other chemical control practices.
- Where possible, make use of predictive disease models to effectively time fungicide applications.
- Where possible, make use of predictive disease models to effectively time fungicide applications.
- Monitor treated fungal populations for resistance development. Notify Nippon Soda Co. Ltd if reduced sensitivity of the pathogen to RENOVO Fungicide is suspected.
- If disease continues to progress after treatment with this product, do not increase the use rate. Discontinue use of this product, and switch to another fungicide with a different target site of action, if available.
- Contact your local extension specialist or certified crop advisor for any additional pesticide resistance-management and/or IPM recommendations for specific crops and pathogens.

For further information and to report suspected resistance, contact Nisso America at (212) 490-0350.

Since the occurrence of resistance cannot be foreseen, Nippon Soda Co., Ltd. accepts no responsibility for any loss of, or damage to, crops resulting from the failure of RENOVO Fungicide to control the resistant fungi strains.

MIXING:

SHAKE WELL BEFORE USING. Some settling may occur during prolonged periods of non-use. When high pH tank mixes exist, the interval between mixing and application should be minimized. The buffering of tank water to pH 6-7 prior to the addition of RENOVO Fungicide is recommended. Add required volume amount of RENOVO Fungicide to partially fill tank (1/2 total volume), agitate by mechanical or hydraulic means, add tank mix product if used (see below), agitate again and then add remaining required amount of water. Continuous agitation is recommended to keep the material in proper suspension. For best results, use spray mixture the same day it is prepared.

This product may be tank mixed with a fertilizer, a supplement, or with registered pest control products, whose labels also allow tank mixing, provided the entirety of both labels, including Directions For Use, Precautions, Restrictions, Environmental Precautions, and Spray Buffer Zones are followed for each product. In cases where these requirements differ between the tank mix partner labels, the most restrictive label must be followed. Do not tank mix products containing the same active ingredient unless specifically listed on this label.

In some cases, tank mixing pest control products can result in reduced pesticide efficacy or increased host crop injury. The user should contact Nisso America at (212) 490-0350 for information before applying any tank mix that is not specifically recommended on this label.

Tank mixing instructions: RENOVO is compatible with most commonly used pesticides. If tank mixing with other materials, add products in the following order: wettable powders, dry flowables, liquid flowables, emulsifiable concentrates, and soluble materials such as fertilizers. No claim of compatibility with other products is implied: Do not tank mix with copper-containing materials or with highly alkaline pesticides, such as Bordeaux mixture or lime sulfur. RENOVO may be applied in conjunction with chemically neutral liquid fertilizers. Application in conjunction with highly alkaline fertilizers, such as aqueous ammonia, may cause a degradation of the pesticide, resulting in reduced performance and should be avoided.

RATE:

On turf, use at least 8 litres of water per 100 m² of area to be sprayed for dollar spot, brown patch and pink snow. Use the recommended rate and reduce spray intervals under conditions of severe disease pressure.

Compatibility:

RENOVO Fungicide is compatible with most pesticides; do not mix with lime or other alkaline materials.

DIRECTIONS FOR USE

As this product is not registered for the control of pests in aquatic systems, DO NOT use to control aquatic pests. DO NOT contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes.

To protect pollinators, follow the instructions regarding bees in the Environmental Precautions section.

Field sprayer application: DO NOT apply during periods of dead calm. Avoid application of this product when winds are gusty. DO NOT apply with spray droplets smaller than the American Society of Agricultural Engineers (ASAE S572.1) medium classification. Boom height must be 60 cm or less above the crop or ground.

Spray Buffer zones:

A spray buffer zone is NOT required for:

- uses with hand-held application equipment permitted on this label.

The spray buffer zones specified in the table below are required between the point of direct application and the closest downwind edge of sensitive terrestrial habitats (such as grasslands, forested areas, shelter belts, woodlots, hedgerows, riparian areas and shrublands), sensitive freshwater habitats (such as lakes, rivers, sloughs, ponds, prairie potholes, creeks, marshes, streams, reservoirs and wetlands) and estuarine/marine habitats.

Method of application	Crop	Spray Buffer Zones (metres) Required for the Protection of:			
		Freshwater Habitat of Depths:		Estuarine/Marine Habitat of Depths:	
		Less than 1 m	Greater than 1 m	Less than 1 m	Greater than 1 m
Field Sprayer	Turf	15	10	1	0

For tank mixes, consult the labels of the tank-mix partners and observe the largest (most restrictive) spray buffer zone of the products involved in the tank mixture and apply using the coarsest spray (ASAE) category indicated on the labels for those tank mix partners.

The spray buffer zones for this product can be modified based on weather conditions and spray equipment configuration by accessing the Buffer Zone Calculator on the Pest Management Regulatory Agency web site.

TURF:

RENOVO Fungicide can be applied to golf course greens, tees, fairways and sod farms only. DO NOT apply to turf in other residential areas including lawns, gardens, parks, playing fields, cemeteries and schools. For use on all fine turf applications of cool and warm season grasses (such as Bentgrass, Blue grass, Fescues, Rye Grass) or their mixtures. RENOVO Fungicide is not phytotoxic to any of the above-mentioned grasses when used in accordance with the label. Apply after mowing or avoid mowing for twelve hours after application. Begin applications, using ground application equipment, when diseases first appear. When disease pressure is severe, apply at the higher recommended rate. Do not exceed 20.65 kg ai/ha per year. DO NOT APPLY BY AIR.

Disease	Rate of Product (mL/100 m ²)	Rate of Active Ingredient (kg ai/ha) ^a	Spray Volume (per 100 m ²)	Spray Schedule	Maximum Seasonal Rate (kg ai/ha)
Dollar Spot (<i>Sclerotinia homoeocarpa</i>)	35 – 42	1.75 – 2.10	Minimum 8 L	Apply when diseases first appear. A second application may be made 10-14 days later if required.	4.20
Brown Patch (<i>Rhizoctonia solani</i>)	42 – 84	2.10 – 4.20	Minimum 8 L	Apply when diseases first appear. If applying at the minimum rate, a second application may be made if required. Allow a minimum interval of 7 days.	4.20
Pink Snow Mould (<i>Microdochium nivale</i>)	175 – 245	8.75 – 12.25	Minimum 8 L	Apply once per season in the late fall, before snowfall for the control of pink snow mould.	12.25

^a Based on 10 000 m²/ha, 1 L/1000mL, and the guarantee of 0.500 kg ai/L.